arXiv:1603.01120v1 [math.CV] 3 Mar 2016

On some subclasses of m-fold symmetric bi-univalent functions

Şahsene Altınkaya* and Sibel Yalçın[1]
*,[1]Department of Mathematics, Faculty of Arts and Science,
Uludag University, Bursa, Turkey.
*corresponding author e-mail: sahsene@uludag.edu.tr
[1]syalcin@uludag.edu.tr

March 4, 2016

Abstract

In this work, we introduce two new subclasses $S_{\Sigma_m}(\alpha,\lambda)$ and $S_{\Sigma_m}(\beta,\lambda)$ of Σ_m consisting of analytic and m-fold symmetric bi-univalent functions in the open unit disk U. Furthermore, for functions in each of the subclasses introduced in this paper, we obtain the coefficient bounds for $|a_{m+1}|$ and $|a_{2m+1}|$.

Keywords: Analytic functions, m-fold symmetric bi-univalent functions, Coefficient bounds.

2010, Mathematics Subject Classification: 30C45, 30C50.

1 Introduction

Let A denote the class of functions of the form

$$f(z)=z+\sum_{n=2}^{\infty}a_nz^n \tag{1}$$

which are analytic in the open unit disk $U=\{z:|z|<1\}$, and let S be the subclass of A consisting of the form (1) which are also univalent in U.

The Koebe one-quarter theorem [7] states that the image of U under every function f from S contains a disk of radius $\frac{1}{4}$. Thus every such univalent function has an inverse f^{-1} which satisfies

$$f^{-1}\left(f\left(z\right)\right)=z \quad (z\in U)$$

and

$$f\left(f^{-1}\left(w\right)\right)=w \quad \left(|w|<r_0\left(f\right)\ ,\ r_0\left(f\right)\geq\frac{1}{4}\right),$$

where

$$f^{-1}(w) = w - a_2w^2 + \left(2a_2^2 - a_3\right)w^3 - \left(5a_2^3 - 5a_2a_3 + a_4\right)w^4 + \cdots . \quad (2)$$

A function $f \in A$ is said to be bi-univalent in U if both f and f^{-1} are univalent in U. Let Σ denote the class of bi-univalent functions defined in the unit disk U.

For a brief history and interesting examples in the class Σ, see [15]. Examples of functions in the class Σ are

$$\frac{z}{1-z}, \quad -\log(1-z), \quad \frac{1}{2}\log\left(\frac{1+z}{1-z}\right)$$

and so on. However, the familier Koebe function is not a member of Σ. Other common examples of functions in S such as

$$z - \frac{z^2}{2} \text{ and } \frac{z}{1-z^2}$$

are also not members of Σ (see [15]).

For each function $f \in S$, the function

$$h(z) = \sqrt[m]{f(z^m)} \qquad (z \in U, \quad m \in \mathbb{N}) \quad (3)$$

is univalent and maps the unit disk U into a region with m-fold symmetry. A function is said to be m-fold symmetric (see [10], [14]) if it has the following normalized form:

$$f(z) = z + \sum_{k=1}^{\infty} a_{mk+1} z^{mk+1} \qquad (z \in U, \quad m \in \mathbb{N}). \quad (4)$$

We denote by S_m the class of m-fold symmetric univalent functions in U, which are normalized by the series expansion (4). In fact, the functions in the class S are *one*-fold symmetric.

Analogous to the concept of m-fold symmetric univalent functions, we here introduced the concept of m-fold symmetric bi-univalent functions. Each function $f \in \Sigma$ generates an m-fold symmetric bi-univalent function for each integer $m \in \mathbb{N}$. The normalized form of f is given as in (4) and the series expansion for f^{-1}, which has been recently proven by Srivastava et al. [17], is given as follows:

$$\begin{aligned} g(w) &= w - a_{m+1}w^{m+1} + \left[(m+1)a_{m+1}^2 - a_{2m+1}\right]w^{2m+1} \quad (5) \\ &\quad - \left[\frac{1}{2}(m+1)(3m+2)a_{m+1}^3 - (3m+2)a_{m+1}a_{2m+1} + a_{3m+1}\right]w^{3m+1} + \cdots . \end{aligned}$$

where $f^{-1} = g$. We denote by Σ_m the class of m-fold symmetric bi-univalent functions in U. For $m = 1$, the formula (5) coincides with the formula (2) of the

class Σ. Some examples of m-fold symmetric bi-univalent functions are given as follows:

$$\left(\frac{z^m}{1-z^m}\right)^m, \quad [-\log(1-z^m)]^{\frac{1}{m}}, \quad \left[\frac{1}{2}\log\left(\frac{1+z^m}{1-z^m}\right)^{\frac{1}{m}}\right].$$

Brannan and Taha [5] introduced certain subclasses of the bi-univalent function class Σ similar to the familiar subclasses. $S^{\star}(\beta)$ and $K(\beta)$ of starlike and convex function of order β $(0 \leq \beta < 1)$ respectively (see [13]). The classes $S_{\Sigma}^{\star}(\alpha)$ and $K_{\Sigma}(\alpha)$ of bi-starlike functions of order α and bi-convex functions of order α, corresponding to the function classes $S^{\star}(\alpha)$ and $K(\alpha)$, were also introduced analogously. For each of the function classes $S_{\Sigma}^{\star}(\alpha)$ and $K_{\Sigma}(\alpha)$, they found non-sharp estimates on the initial coefficients. In fact, the aforecited work of Srivastava et al. [15] essentially revived the investigation of various subclasses of the bi-univalent function class Σ in recent years. Recently, many authors investigated bounds for various subclasses of bi-univalent functions ([1], [2], [8], [11], [15], [16],). Not much is known about the bounds on the general coefficient $|a_n|$ for $n \geq 4$. In the literature, there are only a few works determining the general coefficient bounds $|a_n|$ for the analytic bi-univalent functions ([3], [6], [9]). The coefficient estimate problem for each of $|a_n|$ ($n \in \mathbb{N}\setminus\{1,2\}$; $\mathbb{N} = \{1,2,3,...\}$) is still an open problem.

The aim of the this paper is to introduce two new subclasses of the function class Σ_m and derive estimates on the initial coefficients $|a_{m+1}|$ and $|a_{2m+1}|$ for functions in these new subclasses. We have to remember here the following lemma here so as to derive our basic results:

Lemma 1 *[14] If $p(z) = 1 + p_1 z + p_2 z^2 + p_3 z^3 + \cdots$ is an analytic function in U with positive real part, then*

$$|p_n| \leq 2 \qquad (n \in \mathbb{N} = \{1,2,\ldots\})$$

and

$$\left|p_2 - \frac{p_1^2}{2}\right| \leq 2 - \frac{|p_1|^2}{2}.$$

2 Coefficient bounds for the function class $S_{\Sigma_m}(\alpha,\lambda)$

Definition 2 *A function $f \in \Sigma_m$ is said to be in the class $S_{\Sigma_m}(\alpha,\lambda)$ if the following conditions are satisfied:*

$$\left|\arg\left[\frac{1}{2}\left(\frac{zf'(z)}{f(z)} + \left(\frac{zf'(z)}{f(z)}\right)^{\frac{1}{\lambda}}\right)\right]\right| < \frac{\alpha\pi}{2} \quad (0 < \alpha \leq 1,\ 0 < \lambda \leq 1, \quad z \in U)$$

and

$$\left|\arg\left[\frac{1}{2}\left(\frac{wg'(w)}{g(w)} + \left(\frac{wg'(w)}{g(w)}\right)^{\frac{1}{\lambda}}\right)\right]\right| < \frac{\alpha\pi}{2} \quad (0 < \alpha \leq 1,\ 0 < \lambda \leq 1,\ w \in U)$$

where the function $g = f^{-1}$.

Theorem 3 *Let f given by (4) be in the class $S_{\Sigma_m}(\alpha,\lambda)$, $0<\alpha\leq 1$. Then*

$$|a_{m+1}|\leq \frac{4\lambda\alpha}{m\sqrt{(1+\lambda)\left[4\lambda\alpha+(1+\lambda)(1-\alpha)\right]+2\alpha(1-\lambda)}}$$

and

$$|a_{2m+1}|\leq \frac{2\lambda\alpha}{m\,(1+\lambda)}+\frac{8(m+1)\lambda^2\alpha^2}{m^2(1+\lambda)^2}.$$

Proof. Let $f\in S_{\Sigma_m}(\alpha,\lambda)$. Then

$$\frac{1}{2}\left(\frac{zf'(z)}{f(z)}+\left(\frac{zf'(z)}{f(z)}\right)^{\frac{1}{\lambda}}\right)=[p(z)]^{\alpha} \tag{6}$$

$$\frac{1}{2}\left(\frac{wg'(w)}{g(w)}+\left(\frac{wg'(w)}{g(w)}\right)^{\frac{1}{\lambda}}\right)=[q(w)]^{\alpha} \tag{7}$$

where $g=f^{-1}$, p,q in P and have the forms

$$p(z)=1+p_mz^m+p_{2m}z^{2m}+\cdots$$

and

$$q(w)=1+q_mw^m+q_{2m}w^{2m}+\cdots.$$

Now, equating the coefficients in (6) and (7), we get

$$\frac{m(1+\lambda)}{2\lambda}a_{m+1}=\alpha p_m, \tag{8}$$

$$\frac{m(1+\lambda)}{2\lambda}\left(2a_{2m+1}-a_{m+1}^2\right)+\frac{m^2(1-\lambda)}{4\lambda^2}a_{m+1}^2=\alpha p_{2m}+\tfrac{\alpha(\alpha-1)}{2}p_m^2, \tag{9}$$

and

$$-\frac{m(1+\lambda)}{2\lambda}a_{m+1}=\alpha q_m, \tag{10}$$

$$\frac{m(1+\lambda)}{2\lambda}\left[(2m+1)a_{m+1}^2-2a_{2m+1}\right]+\frac{m^2(1-\lambda)}{4\lambda^2}a_{m+1}^2=\alpha q_{2m}+\tfrac{\alpha(\alpha-1)}{2}q_m^2. \tag{11}$$

From (8) and (10) we obtain

$$p_m=-q_m. \tag{12}$$

and

$$\frac{m^2(1+\lambda)^2}{2\lambda^2}a_{m+1}^2=\alpha^2(p_m^2+q_m^2). \tag{13}$$

Also form (9), (11) and (13) we have

$$\left[\tfrac{m^2(1+\lambda)}{\lambda}+\tfrac{m^2(1-\lambda)}{2\lambda^2}\right]a_{m+1}^2=\alpha\left(p_{2m}+q_{2m}\right)+\tfrac{\alpha(\alpha-1)}{2}(p_m^2+q_m^2).$$

$$=\alpha\left(p_{2m}+q_{2m}\right)+\tfrac{\alpha(\alpha-1)}{2}\tfrac{m^2(1+\lambda)^2}{2\lambda^2\alpha^2}a_{m+1}^2.$$

Therefore, we have

$$a_{m+1}^2 = \frac{4\lambda^2\alpha^2\left(p_{2m}+q_{2m}\right)}{m^2\left\{(1+\lambda)\left[4\lambda\alpha+(1+\lambda)(1-\alpha)\right]+2\alpha(1-\lambda)\right\}}. \tag{14}$$

Appying Lemma 1 for the coefficients p_{2m} and q_{2m}, we obtain

$$|a_{m+1}| \leq \frac{4\lambda\alpha}{m\sqrt{(1+\lambda)\left[4\lambda\alpha+(1+\lambda)(1-\alpha)\right]+2\alpha(1-\lambda)}}.$$

Next, in order to find the bound on $|a_{2m+1}|$, by subtracting (11) from (9), we obtain

$$\frac{2m(1+\lambda)}{\lambda}a_{2m+1} - \frac{m(m+1)(1+\lambda)}{\lambda}a_{m+1}^2 = \alpha\left(p_{2m}-q_{2m}\right) + \tfrac{\alpha(\alpha-1)}{2}(p_m^2-q_m^2).$$

Then, in view of (12) and (13) , and appying Lemma 1 for the coefficients p_{2m}, p_m and q_{2m}, q_m ,we have

$$|a_{2m+1}| \leq \frac{2\lambda\alpha}{m\left(1+\lambda\right)} + \frac{8(m+1)\lambda^2\alpha^2}{m^2(1+\lambda)^2}.$$

This completes the proof of Theorem 3. ■

3 Coefficient bounds for the function class $S_{\Sigma_m}(\beta,\lambda)$

Definition 4 *A function $f \in \Sigma_m$ given by (4) is said to be in the class $S_{\Sigma_m}(\beta,\lambda)$ if the following conditions are satisfied:*

$$\operatorname{Re}\left\{\frac{1}{2}\left(\frac{zf'(z)}{f(z)}+\left(\frac{zf'(z)}{f(z)}\right)^{\frac{1}{\lambda}}\right)\right\} > \beta, \qquad (0\leq\beta<1, \quad 0<\lambda\leq 1, \quad z\in U) \tag{15}$$

and

$$\operatorname{Re}\left\{\frac{1}{2}\left(\frac{wg'(w)}{g(w)}+\left(\frac{wg'(w)}{g(w)}\right)^{\frac{1}{\lambda}}\right)\right\} > \beta, \qquad (0\leq\beta<1, \quad 0<\lambda\leq 1, \quad w\in U). \tag{16}$$

where the function $g = f^{-1}$.

Theorem 5 *Let f given by (4) be in the class $S_{\Sigma_m}(\beta,\lambda)$, $0\leq\beta<1$. Then*

$$|a_{m+1}| \leq \frac{2\lambda}{m}\sqrt{\frac{2\left(1-\beta\right)}{2\lambda^2+\lambda+1}}$$

and

$$|a_{2m+1}| \leq \frac{8(m+1)\lambda^2\left(1-\beta\right)^2}{m^2(1+\lambda)^2} + \frac{2\lambda\left(1-\beta\right)}{m\left(1+\lambda\right)}.$$

Proof. Let $f \in S_{\Sigma_m}(\beta, \lambda)$. Then

$$\frac{1}{2}\left(\frac{zf'(z)}{f(z)} + \left(\frac{zf'(z)}{f(z)}\right)^{\frac{1}{\lambda}}\right) = \beta + (1-\beta)p(z) \tag{17}$$

$$\frac{1}{2}\left(\frac{wg'(w)}{g(w)} + \left(\frac{wg'(w)}{g(w)}\right)^{\frac{1}{\lambda}}\right) = \beta + (1-\beta)q(w) \tag{18}$$

where $p, q \in P$ and $g = f^{-1}$.

It follows from (17) and (18) that

$$\frac{m(1+\lambda)}{2\lambda}a_{m+1} = (1-\beta)p_m, \tag{19}$$

$$\frac{m(1+\lambda)}{2\lambda}\left(2a_{2m+1} - a_{m+1}^2\right) + \frac{m^2(1-\lambda)}{4\lambda^2}a_{m+1}^2 = (1-\beta)p_{2m}, \tag{20}$$

and

$$-\frac{m(1+\lambda)}{2\lambda}a_{m+1} = (1-\beta)q_m, \tag{21}$$

$$\frac{m(1+\lambda)}{2\lambda}\left[(2m+1)a_{m+1}^2 - 2a_{2m+1}\right] + \frac{m^2(1-\lambda)}{4\lambda^2}a_{m+1}^2 = (1-\beta)q_{2m}. \tag{22}$$

From (19) and (21) we obtain

$$p_m = -q_m. \tag{23}$$

and

$$\frac{m^2(1+\lambda)^2}{2\lambda^2}a_{m+1}^2 = (1-\beta)^2(p_m^2 + q_m^2). \tag{24}$$

Adding (20) and (22), we have

$$\left[\frac{m^2(1+\lambda)}{\lambda} + \frac{m^2(1-\lambda)}{2\lambda^2}\right]a_{m+1}^2 = (1-\beta)(p_{2m} + q_{2m}).$$

Therefore, we obtain

$$a_{m+1}^2 = \frac{2\lambda^2(1-\beta)(p_{2m} + q_{2m})}{m^2(2\lambda^2 + \lambda + 1)}.$$

Appying Lemma 1 for the coefficients p_{2m} and q_{2m}, we obtain

$$|a_{m+1}| \leq \frac{2\lambda}{m}\sqrt{\frac{2(1-\beta)}{2\lambda^2 + \lambda + 1}}.$$

Next, in order to find the bound on $|a_{2m+1}|$, by subtracting (22) from (20), we obtain

$$\frac{2m(1+\lambda)}{\lambda}a_{2m+1} - \frac{m(m+1)(1+\lambda)}{\lambda}a_{m+1}^2 = (1-\beta)(p_{2m} - q_{2m}).$$

Then, in view of (23) and (24) , appying Lemma 1 for the coefficients p_{2m}, p_m and q_{2m}, q_m, we have

$$|a_{2m+1}| \leq \frac{8(m+1)\lambda^2 (1-\beta)^2}{m^2(1+\lambda)^2} + \frac{2\lambda (1-\beta)}{m (1+\lambda)}.$$

This completes the proof of Theorem 5. ■

If we set $\lambda = 1$ in Theorems 3 and 5, then the classes $S_{\Sigma_m}(\alpha, \lambda)$ and $S_{\Sigma_m}(\beta, \lambda)$ reduce to the classes $S^{\alpha}_{\Sigma_m}$ and $S^{\beta}_{\Sigma_m}$ and thus, we obtain following corollaries:

Corollary 6 *(see [4]) Let f given by (4) be in the class $S^{\alpha}_{\Sigma_m}$ $(0 < \alpha \leq 1)$. Then*

$$|a_{m+1}| \leq \frac{2\alpha}{m\sqrt{\alpha+1}}$$

and

$$|a_{2m+1}| \leq \frac{\alpha}{m} + \frac{2(m+1)\alpha^2}{m^2}.$$

Corollary 7 *(see [4]) Let f given by (4) be in the class $S^{\beta}_{\Sigma_m}$ $(0 \leq \beta < 1)$. Then*

$$|a_{m+1}| \leq \frac{\sqrt{2(1-\beta)}}{m}$$

and

$$|a_{2m+1}| \leq \frac{2(m+1)(1-\beta)^2}{m^2} + \frac{1-\beta}{m}.$$

The classes $S^{\alpha}_{\Sigma_m}$ and $S^{\beta}_{\Sigma_m}$ are respectively defined as follows:

Definition 8 *(see [4]) A function $f \in \Sigma_m$ given by (4) is said to be in the class $S^{\alpha}_{\Sigma_m}$ if the following conditions are satisfied:*

$$f \in \Sigma, \quad \left|\arg\left(\frac{zf'(z)}{f(z)}\right)\right| < \frac{\alpha\pi}{2} \quad (0 < \alpha \leq 1,\ z \in U)$$

and

$$\left|\arg\left(\frac{wg'(w)}{g(w)}\right)\right| < \frac{\alpha\pi}{2} \quad (0 < \alpha \leq 1,\ w \in U)$$

where the function $g = f^{-1}$.

Definition 9 *(see [4]) A function $f \in \Sigma_m$ given by (4) is said to be in the class $S^{\beta}_{\Sigma_m}$ if the following conditions are satisfied:*

$$f \in \Sigma, \quad \mathrm{Re}\left(\frac{zf'(z)}{f(z)}\right) > \beta \quad (0 \leq \beta < 1, \quad z \in U)$$

and

$$\mathrm{Re}\left(\frac{wg'(w)}{g(w)}\right) > \beta \quad (0 \leq \beta < 1, \quad w \in U)$$

where the function $g = f^{-1}$.

For *one*-fold symmetric bi-univalent functions and $\lambda = 1$, Theorem 3 and Theorem 5 reduce to Corollary 10 and Corollary 11, respectively, which were proven earlier by Murugunsundaramoorthy et al. [12].

Corollary 10 *Let f given by (4) be in the class $S_{\Sigma}^{*}(\alpha)$ $(0 < \alpha \leq 1)$. Then*

$$|a_2| \leq \frac{2\alpha}{\sqrt{\alpha + 1}}$$

and

$$|a_3| \leq 4\alpha^2 + \alpha.$$

Corollary 11 *Let f given by (4) be in the class $S_{\Sigma}^{*}(\beta)$ $(0 \leq \beta < 1)$. Then*

$$|a_2| \leq \sqrt{2\,(1 - \beta)}$$

and

$$|a_3| \leq 4(1 - \beta)^2 + (1 - \beta).$$

References

[1] Altınkaya, Ş., Yalçın, S.: Initial coefficient bounds for a general class of bi-univalent functions, International Journal of Analysis, Article ID 867871, 4 pp, (2014).

[2] Altınkaya, Ş., Yalçın, S.: Coefficient Estimates for Two New Subclasses of Bi-univalent Functions with respect to Symmetric Points, Journal of Function Spaces, Article ID 145242,5 pp, (2015).

[3] Altınkaya, Ş., Yalçın, S.: Faber polynomial coefficient bounds for a subclass ofbi-univalent functions, C.R. Acad. Sci. Paris, Ser.I, 353 (12), 1075–1080, (2015).

[4] Altınkaya, Ş., Yalçın, S.: Coefficient bounds for certain subclasses of m-fold symmetric bi-univalent functions, Journal of Mathematics, Article ID 241683, 5 pp, (2015).

[5] Brannan, D.A., Taha, T.S.: On some classes of bi-univalent functions, Studia Universitatis Babeş-Bolyai Mathematica, 31 (2), 70-77, (1986).

[6] Bulut, S.: Faber polynomial coefficient estimates for a comprehensive subclass of analytic bi-univalent functions, C. R. Acad. Sci. Paris, Ser. I, 352 (6), 479-484, (2014).

[7] Duren, P.L.: Univalent Functions, Grundlehren der Mathematischen Wissenschaften, Springer, New York, USA 259, (1983).

[8] Frasin, B.A., Aouf, M.K.: New subclasses of bi-univalent functions, Applied Mathematics Letters, 24, 1569-1573, (2011).

[9] Hamidi, S.G., Jahangiri, J.M.: Faber polynomial coefficient estimates for analytic bi-close-to-convex functions, C. R. Acad. Sci. Paris, Ser.I, 352 (1), 17–20, (2014).

[10] Koepf, W.: Coefficient of symmetric functions of bounded boundary rotations, Proc. Amer. Math. Soc., 105, 324-329, (1989).

[11] Magesh, N., Yamini, J.: Coefficient bounds for a certain subclass of bi-univalent functions, International Mathematical Forum, 8 (27), 1337-1344, (2013).

[12] Murugusundaramoorty, G., Magesh, N., Prameela, V.: Coefficient bounds for certain classes of bi-univalent function, Abstract and Applied Analysis, Article ID 573017, 3 pp, (2013).

[13] Netanyahu, E.: The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in $|z| < 1$, Archive for Rational Mechanics and Analysis, 32, 100-112, (1969).

[14] Pommerenke, C.: Univalent Functions, Vandenhoeck & Ruprecht, Göttingen, (1975).

[15] Srivastava, H.M., Mishra, A.K., Gochhayat, P.: Certain subclasses of analytic and bi-univalent functions, Applied Mathematics Letters, 23 (10), 1188-1192, (2010).

[16] Srivastava, H.M., Bulut, S., Çağlar, M., Yağmur, N.: Coefficient estimates for a general subclass of analytic and bi-univalent functions, Filomat, 27 (5), 831-842, (2013).

[17] Srivastava, H.M., Sivasubramanian, S., Sivakumar, R.: Initial coefficient boundss for a subclass of m-fold symmetric bi-univalent functions, Tbilisi Math. J., 7 (2), 1-10, (2014).